



UNIVERSITY OF  
**ILLINOIS**  
URBANA - CHAMPAIGN

# **Math 595 Representation-theoretic methods in QIT**

Irreducible representations of the symmetric and unitary groups

---

Felix Leditzky

Fall term 2025

Last update: October 21, 2025

# Conjugacy classes of the symmetric group

---

# Conjugacy classes and irreps

Recall from character theory that for a finite group  $G$ , the number of irreducible representations of  $G$  is equal to the number of conjugacy classes of  $G$ .

Conjugacy classes form a partition of a finite group: The relation  $\sim$  on  $G$  defined as  $g \sim h$  if and only if there exists  $s \in G$  such that  $g = shs^{-1}$  is an equivalence relation.

The *conjugacy classes*  $C_1, \dots, C_k$  of  $G$  are the equivalence classes with respect to this relation, and hence form a partition of  $G$ .

We will need the following facts about permutations (see, e.g., [Goo14]):

## Facts about permutations

- (i) Every permutation  $\pi \in S_n$  can be written uniquely as a product of disjoint cycles, e.g.,  $\pi = (13)(2)(465) \in S_6$ . The *cycle type* of a permutation  $\pi \in S_n$  is the tuple of cycle lengths in non-increasing order. For example,  $\pi = (14)(236)(58)(7)$  has cycle type  $(3, 2, 2, 1)$ .

# Conjugacy classes of the symmetric group

## Facts about permutations

- (ii) Cycle types  $(\lambda_1, \dots, \lambda_d)$  of a permutation  $\pi \in S_n$  form an ordered partition of  $n$ ,  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d \geq 0$  and  $\sum_{i=1}^d \lambda_i = n$ . We use the notation  $\lambda \vdash_d n$  for an ordered partition of  $n$  into at most  $d$  parts. Note: If  $d < n$  then not all possible partitions or cycle types appear.
- (iii) Two permutations  $\pi, \pi' \in S_n$  are conjugate iff they have the same cycle type.

To see (iii), let  $(i_1, \dots, i_k)$  be a cycle of length  $k \leq n$  and  $\sigma \in S_n$  be arbitrary. Then,

$$\sigma(i_1, \dots, i_k)\sigma^{-1} = (\sigma(i_1), \dots, \sigma(i_k)).$$

It follows from (i)-(iii) above that the conjugacy classes of  $S_n$ , and hence its irreducible representations, are indexed by the ordered partitions of  $n$  into  $n$  parts.

# Young diagrams and Young tableaux


---

# Partitions as diagrams


## Definition

Let  $\lambda = (\lambda_1, \dots, \lambda_d) \vdash_d n$  be a partition of  $n$  into at most  $d$  parts. The *Young diagram* corresponding to  $\lambda \vdash_d n$  is an arrangement of  $n$  boxes into  $d$  rows such that the  $i$ -th row has length  $\lambda_i$ .

**Examples:** two Young diagrams  $\lambda_1, \lambda_2 \vdash_4 8$



$$\lambda_1 = (3, 2, 2, 1) \vdash_4 8$$



$$\lambda_2 = (5, 2, 1, 0) \vdash_4 8.$$

# Young tableaux

## Definition (Young tableaux)

- A *Young tableau* is a Young diagram  $\lambda$  with boxes labeled with numbers  $\{1, \dots, N\}$  (we can have  $N \neq n$ ). We call  $\lambda$  the *shape* of the Young tableau  $T$ .
- A *standard Young tableau* is a Young tableau with  $N = n$ , and the labels are strictly increasing along rows (left to right) and along columns (top to bottom).
- A *semistandard Young tableau* is a Young tableau whose labels are non-decreasing along rows and strictly increasing along columns.

The standard Young tableaux of shape  $\lambda = (3, 2)$  are

1	2	3							
4	5								
1	2	4							
3	5								
1	2	5							
3	4								
1	3	4							
2	5								
1	3	5							
2	4								

The semistandard Young tableaux of shape  $\lambda = (3, 2)$  with numbering  $\{1, 2\}$  are

1	1	1							
2	2								
1	1	2							
2	2								

# Indexing irreps in Schur-Weyl decomposition

Schur-Weyl duality gives a decomposition

$$(\mathbb{C}^d)^{\otimes n} = \bigoplus_{\lambda \vdash_d n} V_\lambda \otimes U_\lambda^d.$$

## Irreps in the Schur-Weyl decomposition

In the above decomposition,

- the index  $\lambda \vdash_d n$  runs over the partitions of  $n$  into at most  $d$  parts, or alternatively the set of Young diagrams of  $n$  boxes and at most  $d$  rows;
- the irrep  $V_\lambda$  of  $S_n$  has an orthonormal basis indexed by the set of **standard Young tableaux** of shape  $\lambda \vdash_d n$ .
- the irrep  $U_\lambda^d$  of  $\mathcal{U}_d$  has an orthonormal basis indexed by the set of **semistandard Young tableaux** of shape  $\lambda \vdash_d n$  and numbering  $\{1, \dots, d\}$ .

## Combinatorial formulas for irrep dimensions

Let  $d, n \in \mathbb{N}$ . The number of standard Young tableaux of shape  $\lambda \vdash_d n$  is equal to

$$d_\lambda := \frac{n!}{\prod_{(i,j) \in \lambda} h(i,j)},$$

where for a box  $(i,j)$  in row  $i$  and column  $j$  of  $\lambda$  we define the **hook length**

$$h(i,j) = \#\{\text{boxes to the right of } (i,j)\} + \#\{\text{boxes below } (i,j)\} + \text{the box } (i,j) \text{ itself.}$$

### Example

Let  $\lambda = (4, 2, 1)$  be the Young diagram

6	4	2	1
3	1		
1			

its hook length  $(i,j)$ . Then,

$$d_\lambda = \frac{(4 + 2 + 1)!}{6 \cdot 4 \cdot 2 \cdot 1 \cdot 3 \cdot 1 \cdot 1} = \frac{7!}{6 \cdot 4 \cdot 3 \cdot 2} = 7 \cdot 5 = 35.$$

## Combinatorial formulas for irrep dimensions

The number of semistandard Young tableaux of shape  $\lambda \vdash_d n$  is equal to

$$m_{\lambda,d} = \prod_{1 \leq i < j \leq d} \frac{\lambda_i - \lambda_j + j - i}{j - i}.$$

**Example:**  $d = 3$

The number of semistandard Young tableaux of shape  $\lambda = (4, 2, 1)$  with numbering  $\{1, 2, 3\}$  is

$$m_{\lambda,3} = \frac{4 - 2 + 2 - 1}{2 - 1} \cdot \frac{4 - 1 + 3 - 1}{3 - 1} \cdot \frac{2 - 1 + 3 - 2}{3 - 2} = \frac{3}{1} \cdot \frac{5}{2} \cdot \frac{2}{1} = 15.$$

**Example:**  $d = 4$

The number of semistandard Young tableaux of shape  $\lambda = (4, 2, 1)$  with numbering  $\{1, 2, 3\}$  is

$$m_{\lambda,4} = \frac{4 - 2 + 1}{1} \cdot \frac{4 - 1 + 2}{2} \cdot \frac{4 - 0 + 3}{3} \cdot \frac{2 - 1 + 1}{1} \cdot \frac{2 - 0 + 2}{2} \cdot \frac{1 - 0 + 1}{1} = 3 \cdot \frac{5}{2} \cdot \frac{7}{3} \cdot 2 \cdot 2 \cdot 2 = 140.$$

## Combinatorial formulas for irrep dimensions

The 15 semistandard Young tableaux of shape  $\lambda = (4, 2, 1)$  with numbering  $\{1, 2, 3\}$  are

<table border="1"><tr><td>1</td><td>1</td><td>1</td><td>1</td></tr><tr><td>2</td><td>2</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	1	1	2	2			3				<table border="1"><tr><td>1</td><td>1</td><td>1</td><td>2</td></tr><tr><td>2</td><td>2</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	1	2	2	2			3				<table border="1"><tr><td>1</td><td>1</td><td>1</td><td>3</td></tr><tr><td>2</td><td>2</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	1	3	2	2			3				<table border="1"><tr><td>1</td><td>1</td><td>2</td><td>2</td></tr><tr><td>2</td><td>2</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	2	2	2	2			3				<table border="1"><tr><td>1</td><td>1</td><td>2</td><td>3</td></tr><tr><td>2</td><td>2</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	2	3	2	2			3				<table border="1"><tr><td>1</td><td>1</td><td>3</td><td>3</td></tr><tr><td>2</td><td>2</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	3	3	2	2			3			
1	1	1	1																																																																										
2	2																																																																												
3																																																																													
1	1	1	2																																																																										
2	2																																																																												
3																																																																													
1	1	1	3																																																																										
2	2																																																																												
3																																																																													
1	1	2	2																																																																										
2	2																																																																												
3																																																																													
1	1	2	3																																																																										
2	2																																																																												
3																																																																													
1	1	3	3																																																																										
2	2																																																																												
3																																																																													
<table border="1"><tr><td>1</td><td>1</td><td>1</td><td>1</td></tr><tr><td>2</td><td>3</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	1	1	2	3			3				<table border="1"><tr><td>1</td><td>1</td><td>1</td><td>2</td></tr><tr><td>2</td><td>3</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	1	2	2	3			3				<table border="1"><tr><td>1</td><td>1</td><td>1</td><td>3</td></tr><tr><td>2</td><td>3</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	1	3	2	3			3				<table border="1"><tr><td>1</td><td>1</td><td>2</td><td>2</td></tr><tr><td>2</td><td>3</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	2	2	2	3			3				<table border="1"><tr><td>1</td><td>1</td><td>2</td><td>3</td></tr><tr><td>2</td><td>3</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	2	3	2	3			3				<table border="1"><tr><td>1</td><td>1</td><td>3</td><td>3</td></tr><tr><td>2</td><td>3</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	1	3	3	2	3			3			
1	1	1	1																																																																										
2	3																																																																												
3																																																																													
1	1	1	2																																																																										
2	3																																																																												
3																																																																													
1	1	1	3																																																																										
2	3																																																																												
3																																																																													
1	1	2	2																																																																										
2	3																																																																												
3																																																																													
1	1	2	3																																																																										
2	3																																																																												
3																																																																													
1	1	3	3																																																																										
2	3																																																																												
3																																																																													
<table border="1"><tr><td>1</td><td>2</td><td>2</td><td>2</td></tr><tr><td>2</td><td>3</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	2	2	2	2	3			3				<table border="1"><tr><td>1</td><td>2</td><td>2</td><td>3</td></tr><tr><td>2</td><td>3</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	2	2	3	2	3			3				<table border="1"><tr><td>1</td><td>2</td><td>3</td><td>3</td></tr><tr><td>2</td><td>3</td><td></td><td></td></tr><tr><td>3</td><td></td><td></td><td></td></tr></table>	1	2	3	3	2	3			3																																										
1	2	2	2																																																																										
2	3																																																																												
3																																																																													
1	2	2	3																																																																										
2	3																																																																												
3																																																																													
1	2	3	3																																																																										
2	3																																																																												
3																																																																													

## **Constructing the irreps of the symmetric group**

---

## Row and column group of a Young tableau

Let  $T$  be a standard Young tableau of shape  $\lambda \vdash_d n$ . Define two subgroups  $R_T, C_T$  of  $S_n$  as

$$R_T := \{\pi \in S_n : \pi \text{ permutes integers within rows of } T\}$$

$$C_T := \{\pi \in S_n : \pi \text{ permutes integers within columns of } T\}.$$

### Example

The table below lists standard Young tableau  $T$  of shape  $\lambda \vdash_3 6$  and the corresponding groups  $R_T, C_T$ .

$T$	<table border="1"><tr><td>1</td><td>2</td><td>3</td></tr></table>	1	2	3	<table border="1"><tr><td>1</td><td>2</td></tr><tr><td>3</td><td></td></tr></table>	1	2	3		<table border="1"><tr><td>1</td><td>3</td></tr><tr><td>2</td><td></td></tr></table>	1	3	2		<table border="1"><tr><td>1</td></tr><tr><td>2</td></tr><tr><td>3</td></tr></table>	1	2	3
1	2	3																
1	2																	
3																		
1	3																	
2																		
1																		
2																		
3																		
$R_T$	$S_3$	$\{e, (12)(3)\} \cong S_2$	$\{e, (13)(2)\} \cong S_2$	$\{e\} \cong S_1$														
$C_T$	$\{e\} \cong S_1$	$\{e, (13)(2)\} \cong S_2$	$\{e, (12)(3)\} \cong S_2$	$S_3$														

# Young symmetrizer

We define two elements in  $\mathbb{C}[S_n]$ :

$$r_T := \sum_{\pi \in R_T} \pi$$

$$c_T := \sum_{\pi \in C_T} \operatorname{sgn}(\pi)\pi.$$

## Young symmetrizer

For a given standard Young tableau  $T$  of shape  $\lambda \vdash n$ , the *Young symmetrizer*  $e_T$  is defined as  $e_T := r_T c_T$ .

## Example

- (i) Let  $\lambda = (n) \vdash n$  and  $T = \begin{array}{|c|} \hline 1 \dots n \\ \hline \end{array}$ . Then  $c_T = \{e\}$ ,  $R_T = S_n$  and  $e_T = \sum_{\pi \in S_n} \pi$ .
- (ii) Let  $\lambda = (1, \dots, 1)$  and  $T = \begin{array}{|c|} \hline 1 \\ \vdots \\ n \\ \hline \end{array}$ . Then  $e_T = \sum_{\pi \in S_n} \operatorname{sgn}(\pi)\pi$ .

## Specht modules

Let  $T$  be a Young tableau of shape  $\lambda \vdash n$ , and let  $e_T$  be the corresponding Young symmetrizer. Then  $f_T := \frac{d_\lambda}{n!} e_T$  is the minimal projection in  $\mathbb{C}[S_n]$  corresponding to the irreducible representation  $V_\lambda$  of  $S_n$ , that is,  $V_\lambda \cong \mathbb{C}[S_n]e_T$ . The  $V_\lambda$  are called *Specht modules*. Every irreducible representation of  $S_n$  is isomorphic to a Specht module  $V_\lambda$  for some  $\lambda \vdash n$ , and  $V_\lambda \not\cong V_{\lambda'}$  for  $\lambda \neq \lambda'$ . (Proof: [Chr06] or [Alc18])

Denote by  $\text{SYT}(\lambda)$  the set of all standard Young tableaux of shape  $\lambda$ , and let  $T \in \text{SYT}(\lambda)$ . Then the concrete realization of the group algebra element  $f_T$  on  $(\mathbb{C}^d)^{\otimes n}$  is

$$\Pi_T = \frac{d_\lambda}{n!} \sum_{\pi \in R_T, \sigma \in C_T} \text{sgn}(\sigma) \varphi(\pi) \varphi(\sigma).$$

Recall the isotypical projection  $\Pi_\lambda$  onto the isotypical component  $V_\lambda \otimes U_\lambda^d$  corresponding to the irrep  $V_\lambda$  of  $S_n$ . Then we have the relation

$$\Pi_\lambda = \sum_{T \in \text{SYT}(\lambda)} \Pi_T.$$

# Constructing the irreps of the unitary group

---

Recall the Schur-Weyl decomposition

$$(\mathbb{C}^d)^{\otimes n} \cong \bigoplus_{\lambda \vdash_d n} V_\lambda \otimes U_\lambda^d.$$

We know that the dimension of the  $S_n$ -irrep  $V_\lambda$  is equal to the number  $d_\lambda$  of standard Young tableaux, which can be computed using the hook length formula.

## Irreps of $\mathcal{U}_d$

Let  $\lambda \vdash_d n$  be a Young diagram. Then for each standard Young tableau  $T$  of shape  $\lambda$ , the subspace  $e_T(\mathbb{C}^d)^{\otimes n}$  is an irreducible representation of  $\mathcal{U}_d$  (or equivalently  $GL(\mathbb{C}^d)$ ) isomorphic to  $U_\lambda^d$ .

For an easy proof see [Alc18, Prop. 6.2].

## Constructing a basis for unitary group irreps

Let  $\{|i\rangle\}_{i=1}^d$  be the standard basis of  $\mathbb{C}^d$ . Consider a tensor product basis of  $(\mathbb{C}^d)^{\otimes n}$ :

$$\mathcal{B} = \{|i_1\rangle \otimes \dots \otimes |i_n\rangle : i_j \in [d]\}$$

For a fixed Young tableau  $T$  of shape  $\lambda$  and basis vector  $|v\rangle = |i_1\rangle \otimes \dots \otimes |i_n\rangle \in \mathcal{B}$ , construct a Young tableau  $T_{|v\rangle}$  obtained from replacing  $j$  in  $T$  with the label  $i_j$ .

### Example for $d = 2$ and $n = 4$

$$|v\rangle = |1\rangle \otimes |2\rangle \otimes |1\rangle \otimes |1\rangle : \quad S = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & & \\ \hline \end{array} \quad \longrightarrow \quad S_{|v\rangle} = \begin{array}{|c|c|c|} \hline 1 & 2 & 1 \\ \hline 1 & & \\ \hline \end{array}$$
$$T = \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & & \\ \hline \end{array} \quad \longrightarrow \quad T_{|v\rangle} = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & & \\ \hline \end{array}$$

Note that  $T_{|v\rangle}$  is a semistandard Young tableau, while  $S_{|v\rangle}$  is not.

# Constructing a basis for unitary group irreps

Example for  $d = 2$  and  $n = 4$

$$|v\rangle = |1\rangle \otimes |2\rangle \otimes |1\rangle \otimes |1\rangle: \quad S = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & & \\ \hline \end{array} \longrightarrow S_{|v\rangle} = \begin{array}{|c|c|c|} \hline 1 & 2 & 1 \\ \hline 1 & & \\ \hline \end{array}$$
$$T = \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & & \\ \hline \end{array} \longrightarrow T_{|v\rangle} = \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & & \\ \hline \end{array}.$$

**Observation:** The Young symmetrizer  $e_S$  annihilates the basis vector  $|v\rangle$ , since there is a repetition in  $S_{|v\rangle}$  in the first column, which is *antisymmetrized* by  $e_S$ .

## Corollary

If  $T$  is a Young tableau with more than  $d$  rows, then any basis vector  $|v\rangle \in \mathcal{B}$  is annihilated by  $e_T$  because of the antisymmetrization along columns.

This is why only irreps corresponding to Young diagrams with at most  $d$  rows appear in the Schur-Weyl decomposition.

# Constructing a basis for unitary group irreps

## Generalization:

- (i) For given  $|v\rangle \in \mathcal{B}$  let  $\mu = (\mu_i)_{i=1}^d$  be the vector where  $\mu_i$  is the number of times the basis vector  $|i\rangle$  appears in  $|v\rangle$ .
- (ii) Denote by  $\mu^\downarrow$  the vector obtained from  $\mu$  by sorting its components in non-increasing order. If the standard Young tableau  $T$  has shape  $\lambda$ , then  $e_T$  annihilates  $|v\rangle$  whenever

$$\sum_{i=1}^k \mu_i^\downarrow > \sum_{i=1}^k \lambda_i \quad \text{for some } 1 \leq k \leq d-1.$$

- (iii) This condition is the negation of the **majorization** relation, defined as  $\mu^\downarrow < \lambda$  iff  $\sum_{i=1}^k \mu_i^\downarrow \leq \sum_{i=1}^k \lambda_i$  for all  $k = 1, \dots, d$ .

All basis vectors  $|v\rangle \in \mathcal{B}$  that are not annihilated by  $e_T$  span the irrep  $U_\lambda^d$ , and the ones corresponding to semistandard Young tableaux  $T_{|v\rangle}$  form a basis.

For a concrete way of realizing the irreducible representation of a unitary  $U$  on  $U_\lambda^d$ , we refer to [Mol06] (for an approach using Lie theory) or [FH13, Ex. 15.57] and [Mul07, Sec. 2.1.1] (for a more direct approach).

## Irreps of $S_n$ and $\mathcal{U}_d$

Let  $d = \dim V$  and  $|v\rangle \in V^{\otimes n}$  be non-zero. For a standard Young tableau  $T$  of shape  $\lambda \vdash n$ , consider the Young symmetrizer  $e_T$ . Let  $p$  be the number of parts of the partition  $\lambda$  (or the number of non-zero rows of the Young diagram  $\lambda$ ).

- If  $p \leq d$ , then  $\mathbb{C}[S_n]e_T|v\rangle$  is an irreducible representation of  $S_n$  isomorphic to the Specht module  $V_\lambda$ .
- If  $p \leq d$ , then  $e_T V^{\otimes n}$  is an irreducible representation of  $GL(V)$  (or  $\mathcal{U}_d$ ) on  $V^{\otimes n}$ . These are inequivalent for Young tableaux of different shape.
- Using the above, we have the Schur-Weyl decomposition of  $V^{\otimes n}$  with  $d = \dim V$  as an  $S_n \times \mathcal{U}_d$  representation:

$$V^{\otimes n} = \bigoplus_{\lambda \vdash_d n} V_\lambda \otimes U_\lambda^d.$$

Note that item (iii) of the proposition says that

$$d^n = \sum_{\lambda \vdash_d n} d_\lambda m_{\lambda, d},$$

## **Quantum method of types**

---

# Number of Young diagrams

Fix  $n, d \in \mathbb{N}$  and consider the Schur-Weyl decomposition

$$(\mathbb{C}^d)^{\otimes n} = \bigoplus_{\lambda \vdash_d n} V_\lambda \otimes U_\lambda^d.$$

## Simple counting argument:

Ignoring the constraint of non-decreasing row length ( $\lambda_i \geq \lambda_j$  for  $i \leq j$ ), we can put between 0 and  $n$  boxes in the first row, then in the second row, and so forth, giving the simple (over-)estimate

$$|\{\lambda \vdash_d n\}| \leq (n+1)^d.$$

⇒ only polynomially many terms in Schur-Weyl decomposition.

## Dimension bounds for irreps

Fix  $n, d \in \mathbb{N}$  and consider the Schur-Weyl decomposition

$$(\mathbb{C}^d)^{\otimes n} = \bigoplus_{\lambda \vdash_d n} V_\lambda \otimes U_\lambda^d.$$

For a given Young diagram  $\lambda \vdash_d n$ , define a probability distribution

$$\bar{\lambda} = (\lambda_1/n, \dots, \lambda_d/n).$$

Useful bounds on the dimension of the  $S_n$ -irrep  $V_\lambda$  [Har05]:

$$\exp(nH(\bar{\lambda})) (n+d)^{-d(d+1)/2} \leq \dim V_\lambda \leq \exp(nH(\bar{\lambda})),$$

with **Shannon entropy**  $H(p) = -\sum_i p_i \log p_i$ .

$\Rightarrow$  some  $S_n$ -irreps become **exponentially large** (in  $n$ ).

On the other hand, all  $U_d$ -irreps only grow **polynomially**:

$$\dim U_\lambda^d \leq (n+1)^{d(d-1)/2}.$$

# Permutation-invariant states

A useful consequence of these bounds is the following (see, e.g., [BL25]):

## Permutation-invariant states

Let  $\mathcal{H} = (\mathbb{C}^d)^{\otimes n}$  and  $\rho \in \mathcal{L}(\mathcal{H})$  be a permutation-invariant operator,  $\varphi(\pi)\rho\varphi(\pi)^\dagger = \rho$  for all  $\pi \in S_n$ . Then  $\rho$  is determined by  $\text{poly}(n)$  parameters.

## Proof.

Permutation invariance of  $\rho$  and Schur's Lemma imply that

$$\rho \cong \bigoplus_{\lambda \vdash_d n} \mathbb{1}_{V_\lambda} \otimes \rho_\lambda \quad \text{for some } \rho_\lambda \in \mathcal{L}(U_\lambda^d).$$

Now  $|\{\lambda \vdash_d n\}| \leq (n+1)^d$  and  $\dim U_\lambda^d \leq (n+1)^{d(d-1)/2}$  prove the claim.  $\square$

## Concentration of measure

Denote by  $\Pi_\lambda$  the isotypical projection onto the summand  $V_\lambda \otimes U_\lambda^d$  in the Schur-Weyl decomposition.

Let  $\rho \in \mathcal{L}(\mathbb{C}^d)$  be a quantum state of a  $d$ -dimensional system, and denote by  $s = (s_1, \dots, s_d)$  the spectrum of  $\rho$  ( $s_i \geq s_j$  for  $i \leq j$ ).

Tensor copies  $\rho^{\otimes n}$  are permutation-invariant:  $\rho^{\otimes n} \cong \bigoplus_{\lambda \vdash_{d,n}} \mathbb{1}_{V_\lambda} \otimes \omega_\lambda(\rho)$ .

### Concentration of measure

Using majorization relation and dimension bounds [CM06],

$$\begin{aligned} \exp(-nD(\bar{\lambda} \| s)) (n+d)^{-d(d+1)/2} &\leq \text{tr}(\Pi_\lambda \rho^{\otimes n}) = \dim V_\lambda \text{tr}[\omega_\lambda(\rho)] \\ &\leq \exp(-nD(\bar{\lambda} \| s)) (n+d)^{d(d-1)/2}, \end{aligned}$$

where  $D(p \| q) = \sum_i p_i \log(p_i / q_i)$  is the relative entropy between  $p$  and  $q$ .

The weight of  $\Pi_\lambda \rho^{\otimes n} \Pi_\lambda$  decays exponentially in  $n$  except for those irreps  $\lambda$  for which  $\bar{\lambda}$  is close to the spectrum of  $\rho$  in relative entropy “distance”.

## References

---

- [Alc18] Judith Alcock-Zeilinger. ***The Special Unitary Group, Birdtracks and Applications in QCD***. 2018.
- [BL25] Sujeet Bhalerao and Felix Leditzky. “**Improving quantum communication rates with permutation-invariant codes**”. *arXiv preprint* (Aug. 2025). arXiv: 2508.09978 [quant-ph].
- [Chr06] Matthias Christandl. “**The structure of bipartite quantum states-insights from group theory and cryptography**”. (2006). arXiv: quant-ph/0604183.
- [CM06] Matthias Christandl and Graeme Mitchison. “**The spectra of quantum states and the Kronecker coefficients of the symmetric group**”. *Communications in Mathematical Physics* 261.3 (2006), pp. 789–797. arXiv: quant-ph/0409016 [quant-ph].
- [FH13] William Fulton and Joe Harris. ***Representation theory: a first course***. Vol. 129. Springer Science & Business Media, 2013.
- [Goo14] Frederick Goodman. ***Algebra: Abstract and Concrete, edition 2.6***. SemiSimple Press (Frederick Goodman), 2014. eprint:  
<http://homepage.divms.uiowa.edu/~goodman/algebrabook.dir/book.2.6.pdf>.

- [Har05] Aram W. Harrow. “**Applications of coherent classical communication and the Schur transform to quantum information theory**”. PhD thesis. Massachusetts Institute of Technology, 2005. arXiv: quant-ph/0512255.
- [Mol06] A.I. Molev. “**Gelfand–Tsetlin Bases for Classical Lie Algebras**”. Ed. by M. Hazewinkel. Vol. 4. Handbook of Algebra. North-Holland, 2006, pp. 109–170. arXiv: math/0211289.
- [Mul07] Ketan D. Mulmuley. ***On P vs. NP, Geometric Complexity Theory, and The Flip I: a high-level view***. Tech. rep. TR-2007-13. Computer Science Department: The University of Chicago, 2007. arXiv: 0709.0748 [cs.CC].